

Patient information–infection control department

Endophthalmitis

This leaflet provides information about a rare condition that might occur following an eye operation, procedures or an injury to the eye. Endophthalmitis is treated as an emergency.

What is endophthalmitis?

Endophthalmitis is an inflammation of the internal eye tissues, most commonly caused by an infection.

How will I know that I have endophthalmitis?

If you have recently had an operation on your eye, a procedure such as an intravitreal injection or an injury to your eye and you experience any of the following symptoms, please report immediately to your nearest eye accident and emergency department:

Redness

Sensitivity to light

Vision reduction

Pain

The earlier the eye is treated, the better the recovery.

What are the dangers of endophthalmitis?

The main danger is the risk of eyesight being reduced or lost if treatment is not started as early as possible.

What does the treatment involve?

To find out what is causing endophthalmitis, samples of the fluid from inside your eye will be taken and sent to a laboratory for tests. Antibiotics will be instilled into your eye and antibiotic drops and tablets will be prescribed for you.

If the infection is very severe, you could be admitted into hospital for more intensive treatment such as intravenous antibiotics.

How long will the treatment take?

Your eye will be checked every day and you might need to stay in hospital overnight. When you are discharged, you will be given a date for a follow-up appointment. It is important that you attend this appointment. Once at home, you will need to keep taking the medications prescribed for you (drops

and/or tablets) until the doctor tells you to stop.

Are there any long term side-effects?

Loss of some vision can occur, but blindness is rare.

What will the follow-up procedure involve?

In some cases, treatment could require an operation to remove the infected fluid in the back of the eye.

Subsequent follow-up appointments will depend on the severity of the condition and your response to the treatment. If appropriate, you could be referred to your local eye clinic for follow-up care.

If you require any further advice or information about endophthalmitis, please do not hesitate to contact the infection control team on 020 7253 3411 ext 2539.

Author: Infection control team
Revision number: 2
Approved: September 2014
Review date: September 2016

**Moorfields Eye Hospital NHS
Foundation Trust**
City Road, London EC1V 2PD
Phone: 020 7253 3411
www.moorfields.nhs.uk

Moorfields Direct advice line

Phone: 020 7566 2345

Monday to Friday, 9am to 4.30pm,
for information and advice on eye
conditions and treatments from
experienced ophthalmic-trained nurses.

Patient advice and liaison service (PALS)

Phone: 020 7566 2324 or 020 7566 2325

Email: pals@moorfields.nhs.uk

Moorfields' PALS team provides confidential advice and support to help you sort out any concerns you may have about the care we provide, guiding you through the different services available at Moorfields. The PALS team can also advise you on how to make a complaint.

Your right to treatment within 18 weeks

Under the NHS constitution, all patients have the right to start their consultant-led treatment within 18 weeks of being referred by their GP. Moorfields is committed to fulfilling this right, but if you feel that we have failed to do so, please contact our patient advice and liaison service (PALS) who will be able to advise you further – see above for contact details. For more information about your rights under the NHS constitution, please visit www.nhs.uk/choiceinthenhs.

Moorfields Eye Hospital NHS Foundation Trust
City Road, London EC1V 2PD
Phone: 020 7253 3411
www.moorfields.nhs.uk